

Ocean-Atmosphere State Estimation and Targeted Observing using Coupled Model Ensembles

Craig H. Bishop, PI
Naval Research Laboratory, Marine Meteorology Division
7 Grace Hopper Avenue, Stop 2, Building 702, Room 212
Monterey, CA 93943-5502
Phone: (831) 656 5715, Fax: (831) 656 4769, E-mail: bishop@nrlmry.navy.mil

Dr Sergey Frolov, Visiting Scientist
University Corporation for Atmospheric Research
7 Grace Hopper Avenue, Stop 2, Building 702, Room 212
Monterey, CA 93943-5502
Phone: (831) 656 5715, Fax: (831) 656 4769, E-mail: frolov@nrlmry.navy.mil

Dr. Emanuel Coelho, PI
(local SSC POC: Dr, Charlie Barron, barron@nrlssc.navy.mil), Naval Research Laboratory,
Oceanography Division, Stennis Space Center, Building 1009, Room C128, MS 39529
Phone (Fax): 228 688 5710, E-mail: emanuel.coelho.ctr.po@nrlssc.navy.mil

Dr. James Cummings, Co-PI
Naval Research Laboratory, Marine Meteorology Division
7 Grace Hopper Avenue, Stop 2, Building 702, Room 212
Monterey, CA 93943-5502
Phone: (831) 656 5021, Fax: (831) 656 4769, E-mail: cummings@nrlmry.navy.mil

Dr. Teddy Holt, Co-PI
Naval Research Laboratory, Marine Meteorology Division
7 Grace Hopper Avenue, Stop 2, Building 702, Room 212
Monterey, CA 93943-5502
Phone: (831) 656 4740, Fax: (831) 656 4769, E-mail: holt@nrlmry.navy.mil

Award Number: N0001412WX20323

LONG-TERM GOALS

To build a regional ensemble-based Coupled Ocean Atmosphere Data Assimilation (DA) system that produces probabilistic analyses and forecasts of the ocean-atmosphere state that are significantly superior to those that are currently available. A cornerstone of this endeavour is the assimilation of observations of variables that are affected by both the oceanic and atmospheric state into coupled atmosphere-ocean models such as COAMPS[®]. Examples include scatterometer winds and low-peaking satellite radiance channels that are sensitive to both Sea Surface Temperature (SST) and low-level atmospheric temperature, moisture and aerosol.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2013		2. REPORT TYPE		3. DATES COVERED 00-00-2013 to 00-00-2013	
4. TITLE AND SUBTITLE Ocean-Atmosphere State Estimation and Targeted Observing using Coupled Model Ensembles				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Marine Meteorology Division, 7 Grace Hopper Avenue, Stop 2, Building 702, Room 212, Monterey, CA, 93943-5502				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

OBJECTIVE

Develop a cycling strongly coupled ensemble based data assimilation system for COAMPS[®] that enables ocean state estimates to benefit from observations of atmospheric variables that are influenced by the state of the ocean.

APPROACH

We have built a Hybrid coupled ensemble 3DVAR data assimilation scheme from the mesoscale ocean-atmosphere ensemble forecasting scheme for COAMPS[®] and the 3D-VAR version of the Navy Coupled Ocean Data Assimilation scheme (NCODA), which is currently being used for operations at FNMOC and NAVOCEANO. We have paid particular attention to observations of variables likely to be influenced by the oceanic state such as scatterometer near-ocean-surface winds and other low-level atmospheric observations. Data assimilation is a critical test of an ensemble forecasting system so we also took the time to use information obtained from the data assimilation cycle to improve the quality of the coupled ensemble forecasting system. Because of the high quality of coupled observations associated with it and the on-going Hydrology of the Mediterranean Experiment (HyMeX), we chose to focus on an area encompassing the Mediterranean as our first test region.

WORK COMPLETED IN FY13

- (i) Improved the COAMPS[®] ocean ensemble for the Medeteranian domain by ensuring that the analysis error variance estimates used in the ET were consistent with innovation statistics from profile observations. (Frolov, Bishop, Holt)
- (ii) Built and tested a cycling Hybrid data assimilation scheme that linearly combines a pre-existing static error covariance matrix with a localized ensemble covariance matrix using real observations. (Frolov, Bishop, Holt, Cummings)
- (iii) Built and tested a cycling coupled data assimilation scheme that uses real atmospheric observations to adjust the oceanic state. Currently, the forecast error covariance model for this coupled system is based solely on ensemble covariances because the pre-existing covariance model does not support covariances between atmosphere and ocean (Frolov, Bishop, Holt, Cummings)
- (iv) Devised a perfect model data assimilation test bed in which pseudo-observations are generated from one of the ensemble members and the remaining ensemble members are used to produce a first guess, covariance matrix and an analysis. (Frolov, Bishop, Holt)
- (v) Put together a first draft of a paper demonstrating the remarkable ability of scatterometer winds observations to improve temperature and salinity fields in our perfect model data assimilation experiment. (Frolov, Bishop)

RESULTS

(a) Improvements to the ocean ensemble.

Validation of the legacy ET code for generating ocean ensemble showed that ensembles were grossly under-dispersive (Figure 1a). To constrain the variance of the ET ensemble to be consistent with observations, we used one-month of innovation histories to prescribe profiles of variance. Validation results with the improved ensemble (Figure 1b) show good agreement between ET ensemble variances and temperature innovations.




Figure 1: Ensemble spread diagnostics for ocean temperature

(b) Impact of scatterometer wind observations on ocean temperature and salinity analyses

To demonstrate the potential impact of atmospheric observations on ocean analyses, we devised an identical twin experiment. In the twin experiment, one of 80 coupled COAMPS[®] ensemble members was used to generate pseudo-scatterometer wind observations at every second ocean surface grid point while the other 79 members were used to produce a first guess, covariance matrix and an analysis. We used a simple Gapsari-Cohn horizontal localization with a radius of 200km. No localization was used in the vertical or in-between fluids.

Table 1: RMSE errors for the twin experiment with scatterometer winds.

	$xt-xf$	$xt-xa$	Change
Temperature ($^{\circ}C$)	0.52	0.49	6.3%
Salinity (psu)	0.11	0.10	5.6%
U velocity (m/s)	0.072	0.070	2.7%
V vvelocity (m/s)	0.071	0.069	3.0%

Results of the twin experiment (Table 1) showed that assimilation of scatterometer wind data had a significant projection onto ocean fields and reduced errors in velocity by $\sim 3\%$, temperature by $\sim 6\%$, and salinity fields by $\sim 6\%$.

Closer examination (Figure 2) of the analysis increments showed that spatial patterns of temperature and salinity increments (panels *a* and *c*) had strong similarity to the patterns of forecast errors (panels *b* and *d*).




Figure 2: Surface maps of analysis increments (a, c) and forecast errors (b, d) for the 80-member twin experiment. Arrow in panel a shows the location of profiles plotted in Figure 3. The extent to which the left panels match the right panels shows the extent to which the assimilation step removed error from the state estimate.

We found that the analysis correction from the scatterometer wind observations extended through a considerable depth of the water column (red line in Figure 3). The importance of the spatial coverage of the scatterometer observations was highlighted by computing the correction field that would be obtained if there were only one scatterometer observation located at the horizontal grid point corresponding to the vertical profile of Fig. 3. This single observation correction is shown by the blue line in Fig. 3. Comparison of the blue line with the red line shows that the analysis correction from a single scatterometer wind observation is one or two orders of magnitude smaller than that due to all of the Mediterranean scatterometer observations. This finding demonstrates that much of the potential power of scatterometer wind observations to improve ocean state estimates is associated with the ability of this satellite based instrument to observe a large region of ocean surface winds.

(c) *Effect of atmospheric observations on analysis and forecast skill in cycling system using real observations.*

To further test the impact of atmospheric observations on the ocean analysis, we conducted a preliminary cycling experiment with real-world oceanic (SST and Argo profiles) and atmospheric data (boundary-level temperature and scatterometer winds). In this preliminary experiment we used 20 ensemble members that were localized using Gaspari-Cohn horizontal localization.

Table 2: Forecast errors (y-xf) for a cycling experiment with real data.

	<i>Ocean NCODA</i>	<i>Ocean Ensemble</i>	<i>Coupled Ensemble</i>
<i>SST</i>	0.33	0.41	0.46
<i>Argo Temp</i>	0.30	0.28	0.26
<i>Argo Salt</i>	0.20	0.22	0.21

The results of the cycling DA experiments (Table 2) show that (a) coupled DA had better representation of vertical temperature and salinity structure compared to the ocean-only ensemble; (b) ensemble methods had better representation of the vertical structure of the ocean temperature compared to the existing static NCODA covariance; and (c) ensemble methods had difficulty in fitting sea surface temperature signal.




Figure 3: Profiles of error and increments for location 5W 38N. The black curve gives the correction that would remove all error. The red and blue curves give actual corrections.

These early results are encouraging and we expect further reductions in RMSE scores after we introduce improvements that will include using a larger ensemble size, vertical and inter-fluid localization, and improvements to the generation of the coupled ensemble.

IMPACT/APPLICATIONS

Superior ocean-atmosphere state estimation is the most likely impact of a successful fully coupled ensemble based data assimilation scheme – particularly in data sparse regions. The results of the preliminary experiments described above are very promising in this regard. Better state estimation will lead to better model error diagnoses and better forecasts of the ocean-atmosphere state. Our use of ensemble covariances in coupled data assimilation has and will continue to illuminate inadequacies of the current ensemble analysis system which, in turn, will lead to further improvements of the ensemble system and associated probabilistic forecasts. Better probabilistic forecasts will enable better uncertainty estimation which is critical for the Navy warfighter in interpreting signals from sonic and electromagnetic detection devices. In addition, superior ensemble covariances and localization methods should enable better adaptive sampling methods for environmental state estimation and Warfighter threat detection.

RELATED PROJECTS

“Telescopic 4D State Estimation”. 10/1/12. Office of Naval Research. 3 years.

This project will use ensemble based data assimilation techniques to couple COAMPS atmospheric data assimilation with an ensemble data assimilation scheme for the Navy’s global model.

REFERENCES

Bishop, C.H. and D. Hodyss, 2007: Flow adaptive moderation of spurious ensemble correlations and its use in ensemble based data assimilation. *Quart. J. Roy. Met. Soc.* **133**, 2029-2044.

Bishop, C.H., T.R. Holt, J. Nachamkin, S. Chen, J.G. McLay, J.D. Doyle, and W.T. Thompson, 2009: Regional Ensemble Forecasts Using the Ensemble Transform Technique. *Mon. Wea. Rev.*, **137**, 288–298.